

# West Nile virus encephalitis

By Professor Tom Solomon CBE and Katherine Dodd, University of Liverpool reviewed by Dr James Mitchell, University of Liverpool, and Dr Aravindhan Baheerathan, Imperial College London, UK

## What is West Nile virus encephalitis?

West Nile virus (WNV) encephalitis is an infectious encephalitis caused by West Nile virus. People can get infected from the bite of a mosquito that is infected with the virus. West Nile virus is transferred mainly between pigeons and crows by '*Culex*' mosquitoes. The virus is not transmitted from person to person although there are a small proportion of reported cases where transmission has followed organ transplantation, breast-feeding and blood transfusion. There is no evidence that a person can get the virus from handling live or dead infected birds. However, it is still recommended that barehanded contact with dead animals is avoided.

West Nile fever virus was first discovered in Africa in 1937. Since then, the virus has been found to be distributed not only in Africa, but also southern Europe, the Middle East, America and even parts of the Far East. In 2002 West Nile virus was declared endemic in the USA. The virus has since spread across the North American continent into Canada (where it is now endemic) and the Caribbean, although only very few cases have been reported in Latin America. Recently, outbreaks have also occurred in Europe, and in 2016 included Hungary, Serbia, Austria, Italy, Romania, Ukraine and Spain. Surveillance is currently taking place in the southeast of the UK, however, the virus has not been found in any mosquitos in this country.

## Symptoms of West Nile virus infection/ West Nile virus encephalitis

The time from infection to developing symptoms of West Nile infection is usually 3 - 14 days. The majority (80%) of people who get infected do not display any symptoms. Mild infections are common and include a high temperature, headache and body aches, often with skin rash and swollen lymph glands.

Until recently, brain problems after West Nile infection were thought to be rare. However, in recent years the virus is causing different types of disease affecting the central nervous system including

meningitis (inflammation of the lining of the brain), encephalitis (direct inflammation of the brain) and myelitis (inflammation of the spinal cord). Approximately 1 in 100 people infected with the virus will develop brain complications as a result of infection. Whilst serious illness can occur in people of any age, those at highest risk of severe illness include those aged greater than 50 and those who are immunocompromised (people whose immune system doesn't work as well as it should do).

Encephalitis results when the virus invades the brain causing damage and an inflammatory (swelling) response. The symptoms and signs of West Nile encephalitis range from muscle weakness and paralysis, mild confusion and behavioural changes (which may be mistaken for hysteria) to convulsions (fits) and deep coma.

## Diagnosis and prognosis of West Nile encephalitis

Diagnosis is made based on the symptoms and in most cases can be confirmed by urine, blood or cerebrospinal fluid (fluid surrounding the brain and the spinal cord) tests that detect the virus or the body's immune response to the virus. West Nile virus encephalitis is associated with considerable short-term and long-term risks of death and significant difficulties such as fatigue, weakness, depression, difficulty walking, ataxia (difficulties with coordination, balance and speech) and memory loss.

#### **Treatment and prevention**

There is no specific treatment for the cause of West Nile virus encephalitis and treatment is supportive. Treatment, in most cases, consists of treating the complications of the disease such as high fever and aches, low blood pressure, blood loss, convulsions (fits), or raised pressure within the skull.

The simplest preventative measure is to avoid bites from the mosquitoes that carry the virus. This means wearing long sleeves and trousers, especially during the evening when the mosquitos bite more frequently and avoiding areas where stagnant water can be found as mosquito larvae need still water to develop. For further protection use an insect spray containing at least 30% DEET (N,N-diethyl-3methlybenzamide) and sleep under bed-nets.

There is no vaccine at present to protect against West Nile virus, although several are being developed and tested.

### **Future research**

Current research is looking at ways to reduce resistance to insecticides used to eradicate mosquitos and develop the best possible treatments for the disease. Treatments that have been tried include anti-viral drugs and drugs that modulate the body's immune system. It is hoped that a vaccine will soon be available for use, although work needs to be done to ensure that the effects are long-lasting and safe. Much research is also being carried out on surveillance of the disease and to try to predict when and where outbreaks are going to happen.

FS010V4 West Nile encephalitis

Date created: March 2001/Last updated: March 2022/ Review date: March 2025

<u>Disclaimer</u>: We try to ensure that the information is easy to understand, accurate and up to date as possible. If you would like more information on the source material and references the author used to write this document, please contact Encephalitis International. None of the authors of the above document has declared any conflict of interest, which may arise from being named as an author of this document.

#### Support our information

With our support, no one has to face encephalitis alone. Our advice and information is available free of charge to everyone affected but we are truly grateful when supporters feel able to contribute a little to the cost of these resources. Please make a donation today by visiting <u>www.encephalitis.info/donate</u> or text the word DOCTOR to 70085 to donate £5.

Thank you!

Encephalitis International, 32 Castlegate, Malton, North Yorkshire, YO17 7DT, UK

Administration: +44 (0) 1653 692583 Support: +44 (0) 1653 699599

Email: mail@encephalitis.info Website: www.encephalitis.info

Encephalitis International is the operating name of the Encephalitis Support Group.

Registered Charity England and Wales No: 1087843; Registered Charity in Scotland No: SC048210

Charitable Company registered in England and Wales No: 04189027