

## **Nipah virus infection and encephalitis**

By Alina Ellerington, Encephalitis International and reviewed by Dr Ali Alam, University of Liverpool

### **What is Nipah virus infection?**

Nipah virus infection is a zoonotic disease (a disease that jumps from animals to humans) caused by the Nipah virus. Fruit bats are the natural hosts of the Nipah virus. The virus spreads from bats to pigs and then to humans through close contact with infected pigs or their contaminated tissues. Human-to-human transmission is also possible.

In humans, transmission occurs via consumption of food contaminated with the virus and via contact with infected animals or human body fluids (blood, urine, saliva). The disease was first reported in 1998 in Malaysia with human infections resulting from direct contact with sick pigs or consumption of meat contaminated by the virus. Outbreaks also occurred in Bangladesh, Philippines, Singapore, and India where transmission resulted from consumption of fruits or fruit products (such as raw date palm juice) contaminated with urine or saliva from infected fruit bats. Human to human transmission was seen among families/carers of those affected in hospital settings.

### **Symptoms of Nipah virus encephalitis**

Symptoms of infection range from mild to severe and include fever, difficulty breathing, cough, headache, pneumonia, and vomiting. However, Nipah virus is also neuroinvasive and can cause a devastating, contagious encephalitis outbreak, characterised by severe brain inflammation and dysfunction. Encephalitis due to Nipah virus infection manifests as alteration of the level of consciousness, neuropsychiatric symptoms, seizures or coma. In some cases, encephalitis can appear several months to years (late onset) after recovering from initial infection. It has a high mortality rate (59%).

### **Diagnosis of Nipah virus encephalitis**

Diagnosis can be made during the acute illness by identification of the virus in urine or blood, by nasal and throat swabs or cerebrospinal fluid (CSF), or later in the illness/during recovery by identification of specific antibodies.

### **Treatment**

There isn't a specific antiviral or vaccine against this virus. Treatment is limited to supportive care through rest, hydration, and treatment of specific symptoms as they occur. Medications include acetaminophen and/or ibuprofen to relieve pain and fevers; dimenhydrinate and/or ondansetron to control vomiting; and dexamethasone or similar

inhalers to improve respiratory symptoms. Anti-seizure medications, such as benzodiazepines, may be used to alleviate seizures associated with acute encephalitis and maintain control of neurological symptoms.

### **Outcomes of Nipah virus encephalitis**

Those who survive can suffer a brain injury and can be left with long-term difficulties such as changes in behaviour, eye movement problems, neck muscle spasms, weakness, facial paralysis, depression, memory problems, and fatigue.

### **Prevention**

Prevention measures consist of avoiding exposure to sick pigs and bats in areas where the virus is present, and not drinking raw date palm sap which can be contaminated by an infected bat. During outbreaks, infection prevention and control measures, and personal protective equipment are required. The practice of good hand hygiene cannot be overstated: frequent handwashing with soap and water can help prevent transmission via contact with contaminated surfaces. Finally, it is important to disinfect equipment and surfaces, particularly medical equipment used on Nipah patients.

---

FS072V1 Nipah virus infection and encephalitis

Page created: January 2024; Last update: January 2024; Review date: January 2027

### **Support our information**

With our support, no one has to face encephalitis alone. Our advice and information are available free of charge to everyone affected but we are truly grateful when supporters feel able to contribute a little to the cost of these resources. Please make a donation today by visiting [www.encephalitis.info/donate](http://www.encephalitis.info/donate) or text the word DOCTOR to 70085 to donate £5.

### **Thank you!**

---

Disclaimer: We try to ensure that the information is easy to understand, accurate and up to date as possible. If you would like more information on the source material and references the author used to write this document, please contact Encephalitis International. None of the authors of the above document has declared any conflict of interest, which may arise from being named as an author of this document.

---

**Encephalitis International**, 32 Castlegate, Malton, North Yorkshire, YO17 7DT, UK

**Administration:** +44 (0) 1653 692583 **Support:** +44 (0) 1653 699599

**Email:** [mail@encephalitis.info](mailto:mail@encephalitis.info) **Website:** [www.encephalitis.info](http://www.encephalitis.info)

Encephalitis International is the operating name of the Encephalitis Support Group.

Registered Charity England and Wales No: 1087843; Registered Charity in Scotland No: SC048210

Charitable Company registered in England and Wales No: 04189027